



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

equation of the curve formed by the perpetual intersection of rays, which, diverging from a luminous point, are reflected by a polished surface of a given curvature. Curves of this description have been denominated caustics; and the method usually employed to discover their polar equations, or the relation between the radius vector of any point of the curve and the tangent at that point, is both long and inelegant, and is considered by the author as involving considerable inaccuracy of reasoning. He proposes, therefore, to substitute a new method of investigation, by taking the polar equation of one of the reflected rays, and differentiating this equation with respect to the arbitrary quantities solely which determine its position, and thus obtaining the polar co-ordinates of the point of intersection of two consecutive lines; and finally, by elimination, the equation of the curve in which all such points are found. He is thus led to results remarkable for their simplicity, elegance, and generality: and he gives particular applications of his method, exemplifying the facility with which it effects the solution of problems extremely difficult of management by the ordinary methods hitherto employed. His method is also applicable to the determination of the equations of the evolutes of curves, and to various other problems of a similar nature.

A paper was also read, entitled, "Discovery of the Metamorphoses in the second Type of the Cirripedes, viz. the *Lepadæ*, completing the Natural History of these singular Animals, and confirming their affinity with the Crustacea." By J. V. Thompson, Esq., F.L.S., Deputy Inspector General of Hospitals. Communicated by Sir James Macgrigor, Bart., M.D., F.R.S.

The discoveries made by the author of the remarkable metamorphoses which the animals composing the first family of the Cirripedes, or *Balani*, undergo in the progress of their developement, and which he has published in the third number of his *Zoological Researches* (p. 76), are in the present paper, which is intended as a prize Essay for one of the Royal Medals, followed up by the report of his discovery of similar changes exhibited by three species of two other genera of the second tribe of this family, namely, the *Lepadæ*. The larvæ of this tribe, like those of the *Balani*, have the external appearance of bivalve *Monoculi*, furnished with locomotive organs, in the form of three pairs of members, the most anterior of which are simple and the other bifid. The back of the animal is covered by an ample shield, terminating anteriorly in two extended horns, and posteriorly in a single elongated spinous process. Thus they possess considerable powers of locomotion, which, with the assistance of an organ of vision, enable them to seek their future permanent place of residence. The author is led from his researches to the conclusion that the Cirripedes do not constitute, as modern naturalists have considered them, a distinct class of animals, but that they occupy a place intermediate between the Crustacea decapoda, with which the *Balani* have a marked affinity, and the Crustacea entomostraca, to which the *Lepadæ* are allied; and that they have no natural affinity with the Testaceous Mollusca, as was supposed by Linnæus, and all the older systematic writers on Zoology.